

June 22, 2010

TUM-HEP-757/10

The Impact of a 4th Generation on Mixing and CP Violation in the Charm System

Andrzej J. Buras^{a,b}, Björn Duling^a, Thorsten Feldmann^a,
Tillmann Heidsieck^a, Christoph Promberger^a, Stefan Recksiegel^a

^a*Physik Department, Technische Universität München, James-Frank-Straße,
D-85748 Garching, Germany*

^b*TUM Institute for Advanced Study, Technische Universität München, Arcisstr. 21,
D-80333 München, Germany*

Abstract

We study $D^0 - \bar{D}^0$ mixing in the presence of a fourth generation of quarks. In particular, we calculate the size of the allowed CP violation which is found at the observable level well beyond anything possible with CKM dynamics. We calculate the semileptonic asymmetry $a_{\text{SL}}(D)$ and the mixing induced CP asymmetry $\eta_f S_f(D)$ which are correlated with each other. We also investigate the correlation of $\eta_f S_f(D)$ with a number of prominent observables in other mesonic systems like ε'/ε , $\text{Br}(K_L \rightarrow \pi^0 \nu \bar{\nu})$, $\text{Br}(K^+ \rightarrow \pi^+ \nu \bar{\nu})$, $\text{Br}(B_s \rightarrow \mu^+ \mu^-)$, $\text{Br}(B_d \rightarrow \mu^+ \mu^-)$ and finally $S_{\psi\phi}(B_s)$ in the B_s system. We identify a clear pattern of flavour and CP violation predicted by the SM4 model: While simultaneous large 4G effects in the K and D systems are possible, accompanying large NP effects in the B_d system are disfavoured. However this behaviour is not as pronounced as found for the LHT and RSc models. In contrast to this, sizeable CP violating effects in the B_s system are possible unless extreme effects in $\eta_f S_f(D)$ are found, and $\text{Br}(B_s \rightarrow \mu^+ \mu^-)$ can be strongly enhanced regardless of the situation in the D system. We find that, on the other hand, $S_{\psi\phi}(B_s) > 0.2$ combined with the measured ε'/ε significantly diminishes 4G effects within the D system.

1 Introduction

The addition of a sequential fourth generation (4G) of quarks and leptons to the SM (hereafter referred to as SM3) is undoubtedly one of its simplest extensions. Reviews and summary statements of the thus extended model (hereafter referred to as SM4) can be found in [1, 2]. While not addressing any of the known hierarchy and naturalness problems, if present in nature a 4G is likely to have a number of profound implications as clearly seen in the most recent literature. In particular during the last years, a number of analyses were published with the goal of investigating the impact of the existence of a 4G on Higgs physics [3, 4], electroweak precision tests [3, 5–8], renormalisation group effects [9, 10] and flavour physics [1, 11–21]. Also detailed analyses of supersymmetry in the presence of a 4G have recently been performed in [22, 23], where a good up to date collection of references to papers on the SM4 is contained in the second of these papers.

From the point of view of flavour changing neutral current (FCNC) processes, the SM4 is particularly interesting since, although it introduces non-Minimally Flavour Violating (MFV) interactions, it contains much fewer parameters than other New Physics (NP) scenarios like the Littlest Higgs model with T-parity (LHT), Randall-Sundrum (RS) models or the general MSSM. Moreover, the fact that the operator structure, similarly to the LHT model, is not modified with respect to the SM3, implies that the non-perturbative uncertainties in the SM4 are at the same level as in the SM3. On the other hand, the absence of tree level contributions to FCNC processes, in particular to particle-antiparticle mixing, and the absence of left-right operators allows to satisfy the experimental constraint from the CP violating parameter ε_K more easily than this is possible in RS models. Finally, the non-decoupling effects of the 4G fermions and the fact that the masses of new fermions are accessible easily to the LHC allows us to expect that these NP scenario can be simultaneously tested by high energy and flavour experiments.

In [20] we have performed a detailed analysis of non-MFV effects in the K , B_d and B_s systems in the SM4, paying particular attention to correlations between flavour observables and addressing within this framework a number of anomalies present in the experimental data. Similarly to the LHT model, the RS model with custodial symmetry (RSc) and supersymmetric flavour models which have been analysed in [24–29], we have found that the SM4 can strongly violate correlations between various observables characteristic for models with MFV and in particular with Constrained Minimal Flavour Violation (CMFV). At the same time it is still possible to satisfy all existing data on flavour violating processes and electroweak precision observables.

Probably the most striking signature of the SM4, compared to the LHT, RSc and SUSY flavour models, is the possibility of having simultaneously sizeable NP effects in

the K , B_d and B_s systems, even if truly spectacular effects are only possible in rare K decays. The prominent exception is $S_{\psi\phi}(B_s)$ which can also be enhanced by more than an order of magnitude — although the imposition of the ε'/ε constraint has significant impact on the size of this enhancement. This different pattern can be traced back to the fact that the mass scales involved in the SM4 are generally significantly lower than in the LHT and in particular in the RSc. Also the non-decoupling effects present in the SM4 which are absent in the LHT, RSc and SUSY play a role here.

The main goal of the present paper is the extension of our previous analysis [20] to $D^0 - \bar{D}^0$ mixing and in particular to the mixing-induced CP violation in the charm system, where CP violating effects are predicted to be tiny within the SM3¹. The three basic questions that we want to address are as follows:

- Are large 4G effects in the charm system still possible if consistence with all available constraints from tree level decays, particle-antiparticle mixing, ε'/ε and rare decays in the K , B_d and B_s systems is required and also electroweak precision tests are taken into account?
- Can the large 4G effects in the K , B_d and B_s systems found by us in [20] still be maintained in the presence of constraints from charm data?
- How are the 4G effects in the K , B_d and B_s systems correlated with the corresponding effects in the charm system? In fact, strong correlations between the $D^0 - \bar{D}^0$ and $K^0 - \bar{K}^0$ systems in models with purely left-handed currents have been pointed out in [31] and analysed also in [32].

From the point of view of charm physics, the present paper follows the strategy of our previous papers on this topic in the context of the LHT model [32] and supersymmetric models [33]. We will not repeat in detail the description of the SM4 and of the structure of the 4G mixing matrix since a detailed presentation of this model can be found in [20], where the same notation as in the present paper is used. Similarly, we will not repeat the formalism of $D^0 - \bar{D}^0$ mixing as it is presented in [29, 31, 32, 34–36] in a notation suitable for our purposes.

Our paper is organised as follows. In Section 2 we briefly recapitulate the basic ingredients of the SM4 which are needed for our analysis and we present the effective Hamiltonian for $D^0 - \bar{D}^0$ mixing in this model. Subsequently, we discuss in explicit terms the connection between D and K physics pointed out in [31], which in the SM4 turns out to be not as transparent as in LHT [32]. To this end we consider different scenarios for the V_{4G} mixing matrix which have been analysed in detail in our previous paper [20].

¹See however [30].

Section 3 is devoted to our numerical analysis. Here we study a number of correlations between different observables and we address the three questions posed above. We summarise our results in Section 4.

2 Relevant Formulae and Analytical Considerations

2.1 The Effective Hamiltonian

The effective Hamiltonian for $\Delta C = 2$ transitions following from the usual box diagrams and including the contributions from the b' -quark is given as follows

$$\begin{aligned} \mathcal{H}_{\text{eff}}^{\Delta C=2} = & \frac{G_F^2}{16\pi^2} M_W^2 \left[\lambda_s^{(D)^2} \eta_{ss}^{(D)} S_0(x_s) + \lambda_b^{(D)^2} \eta_{bb}^{(D)} S_0(x_b) + \lambda_{b'}^{(D)^2} \eta_{b'b'}^{(D)} S_0(x_{b'}) \right. \\ & + 2\lambda_b^{(D)} \lambda_s^{(D)} \eta_{bs}^{(D)} S_0(x_b, x_s) + 2\lambda_{b'}^{(D)} \lambda_s^{(D)} \eta_{b's}^{(D)} S_0(x_{b'}, x_s) \\ & \left. + 2\lambda_{b'}^{(D)} \lambda_b^{(D)} \eta_{b'b}^{(D)} S_0(x_{b'}, x_b) \right] \left[\alpha_s^{(4)}(\mu) \right]^{-6/25} \left[1 + \frac{\alpha_s^{(4)}(\mu)}{4\pi} J_4 \right] Q(\Delta C = 2) + h.c. , \end{aligned} \quad (2.1)$$

where

$$Q(\Delta C = 2) = (\bar{u}c)_{V-A}(\bar{u}c)_{V-A} , \quad (2.2)$$

and

$$\lambda_i^{(D)} = V_{ci}^* V_{ui} , \quad (2.3)$$

which satisfy the unitarity relation

$$\lambda_d^{(D)} + \lambda_s^{(D)} + \lambda_b^{(D)} + \lambda_{b'}^{(D)} = 0 . \quad (2.4)$$

For the QCD corrections we will use the approximate relations

$$\eta_{b'b'}^{(D)} \approx \eta_{tt}^{(K)} , \quad \eta_{b'b}^{(D)} \approx \eta_{b's}^{(D)} \approx \eta_{ct}^{(K)} , \quad \eta_{ss}^{(D)} \approx \eta_{bb}^{(D)} \approx \eta_{bs}^{(D)} \approx \eta_{cc}^{(K)} , \quad (2.5)$$

which are found by inspecting the structure of QCD corrections in the $D^0 - \bar{D}^0$ and $K^0 - \bar{K}^0$ systems. In this context let us remark that with $m_{b'}$ defined as $m_{b'}(m_{b'})$ the QCD factors are only weakly dependent on $m_{b'}$. Moreover we recall that the μ -dependent QCD corrections in (2.1) are absorbed in the renormalisation group invariant parameter $\hat{B}_D$. This parameter is defined by

$$\hat{B}_D = B_D(\mu) \left[\alpha_s^{(4)}(\mu) \right]^{-6/25} \left[1 + \frac{\alpha_s^{(4)}(\mu)}{4\pi} J_4 \right] . \quad (2.6)$$

From lattice calculations [37–39], one has $B_D(\mu = 2 \text{ GeV}) = 0.845 \pm 0.024_{-0.006}^{+0.024}$, and with $J_4 = 6719/3750$ [40] and $\alpha_s(M_Z) = 0.1184 \pm 0.0007$ [41], we find

$$\hat{B}_D = 1.18_{-0.05}^{+0.07}, \quad (2.7)$$

which will be used in our numerical calculations.

2.2 CP Violation and Mixing Parameters

The short distance (SD) contributions to the matrix element of the effective Hamiltonian (2.1) can be written as

$$\langle \bar{D}^0 | \mathcal{H}_{\text{eff}}^{\Delta C=2} | D^0 \rangle_{\text{SD}} \equiv |M_{12}^D| e^{2i\phi_D} = (M_{12}^D)^*, \quad (2.8)$$

where

$$M_{12}^D = \frac{G_F^2}{12\pi^2} F_D^2 \hat{B}_D m_D M_W^2 \overline{M}_{12}^D, \quad (2.9)$$

with

$$\begin{aligned} \overline{M}_{12}^D = & \lambda_s^{(D)*2} \eta_{cc}^{(K)} S_0(x_s) + \lambda_b^{(D)*2} \eta_{cc}^{(K)} S_0(x_b) + \lambda_{b'}^{(D)*2} \eta_{tt}^{(K)} S_0(x_{b'}) \\ & + 2\lambda_b^{(D)*} \lambda_s^{(D)*} \eta_{cc}^{(K)} S_0(x_b, x_s) + 2\lambda_{b'}^{(D)*} \lambda_s^{(D)*} \eta_{ct}^{(K)} S_0(x_{b'}, x_s) + 2\lambda_{b'}^{(D)*} \lambda_b^{(D)*} \eta_{ct}^{(K)} S_0(x_{b'}, x_b). \end{aligned} \quad (2.10)$$

Here we used the approximations given in (2.5). The full matrix elements then are given by

$$\langle \bar{D}^0 | \mathcal{H}_{\text{eff}}^{\Delta C=2} | D^0 \rangle = (M_{12}^D + M_{12}^{\text{LD}})^* - \frac{i}{2} \Gamma_{12}^{\text{LD}*}, \quad (2.11)$$

$$\langle D^0 | \mathcal{H}_{\text{eff}}^{\Delta C=2} | \bar{D}^0 \rangle = (M_{12}^D + M_{12}^{\text{LD}}) - \frac{i}{2} \Gamma_{12}^{\text{LD}}. \quad (2.12)$$

Here Γ_{12}^{LD} and M_{12}^{LD} stand for long distance (LD) contributions with the former arising exclusively from SM3 dynamics. These contributions are very difficult to estimate and will be included in our phenomenological analysis using the strategy of [32, 33].

2.3 Connections Between D and K Physics and Beyond

In [31], the connection between $D^0 - \bar{D}^0$ and $K^0 - \bar{K}^0$ mixing has been discussed within the framework of approximately $SU(2)_L$ -invariant NP models. Due to the connection between up- and down-type quarks in the SM3 through the CKM matrix, the NP contributions to $D^0 - \bar{D}^0$ and $K^0 - \bar{K}^0$ mixing are not independent of each other. This

observation has been used in [31] to derive lower bounds on the NP scale in various NP scenarios, which emerge if the experimental constraints on $D^0 - \bar{D}^0$ and $K^0 - \bar{K}^0$ mixing are applied to only the $(V - A) \otimes (V - A)$ contribution.

The LHT model which contains only $(V - A) \otimes (V - A)$ operators belongs to the class of models where the connection between D and K physics is particularly transparent. A detailed analysis in this context has been performed in [32]. There it has been found that

- The correlation between the K and D systems implies that large NP effects in K and D decays are possible simultaneously.
- On the other hand simultaneous large NP effects in D and B decays in the LHT model are unlikely. Indeed while either the CP asymmetry $S_{\psi\phi}(B_s)$ in $B_s - \bar{B}_s$ mixing or $|q/p|$ in the $D^0 - \bar{D}^0$ system can deviate significantly from their SM3 predictions, it is unlikely to observe large deviations from the SM3 in both quantities simultaneously. The improved measurements of $S_{\psi\phi}(B_s)$ at the Tevatron and the LHC in the coming years will therefore have a large impact on the possible size of CP violating effects in D decays within the LHT model.
- The above observations are consistent with the fact that large simultaneous effects in K and B decays in the LHT are found to be very unlikely [25, 26, 42].

It is of interest to understand whether this particular pattern of correlations between $D^0 - \bar{D}^0$ and $K^0 - \bar{K}^0$ mixing can also be found in the SM4. Analysing the structure of the SM4 contributions we find, although the SM4 again only involves purely left-handed operators, that NP contributes in a different manner as compared to the LHT model or the general type of models studied in [31]. The most important difference is the non-unitarity of the 3×3 submatrix of the CKM matrix in the SM4. Therefore not only the short-distance Wilson coefficients of the $(V - A) \otimes (V - A)$ operators are modified, but also the CKM elements in front of the SM3 contributions generally depend on the 4G mixing angles and phases. From this, it is immediately clear that—in contrast to the situation assumed in [31]—the primary interest is not to use $D^0 - \bar{D}^0$ and $K^0 - \bar{K}^0$ mixing observables in order to derive a lower bound on the relevant NP mass scale. In the contrary, after fixing reasonable ranges for the t' and b' masses from considerations in the electroweak sector, the flavour observables now generally depend in a rather complicated manner on the 4G mixing parameters, the short-distance loop functions involving 4G and SM3 quarks, and long-distance matrix elements. Focusing on the $D^0 - \bar{D}^0$ sector and using the standard parametrisation of the 4G mixing matrix together with $\theta_{ij} \ll 1$,

one finds the rather simple approximate expressions

$$\lambda_s^{(D)} \simeq s_{12}, \quad \lambda_b^{(D)} \simeq s_{13} s_{23} e^{-i\delta_{13}}, \quad \lambda_{b'}^{(D)} \simeq \sigma_{12} \equiv s_{14} s_{24} e^{i(\delta_{24} - \delta_{14})}. \quad (2.13)$$

We observe that the new contributions from the b' -quark essentially constrain the magnitude and phase of the parameter combination σ_{12} . In particular, we expect sizable contributions to $D^0 - \bar{D}^0$ mixing observables if $\lambda_{b'}^{(D)}$ competes with $\lambda_b^{(D)}$ appearing in front of the short-distance contribution in the SM3, i.e. if

$$s_{14} s_{24} \gtrsim s_{13} s_{23} \sim \lambda^5, \quad (2.14)$$

where we indicated the scaling with the Wolfenstein parameter $\lambda = s_{12}$. The analogous expressions for the CKM factors $\lambda_t^{(K)}$ and $\lambda_{t'}^{(K)}$, relevant for $K^0 - \bar{K}^0$ mixing, can be found in Section 9.2 of [20]. Here, it is generally found that the expressions depend on all five new 4G mixing parameters. Consequently, the correlation between $D^0 - \bar{D}^0$ and $K^0 - \bar{K}^0$ mixing observables is rather involved. Some simplification arises if one concentrates on specific scenarios, where the 4G mixing angles scale with particular powers of the Wolfenstein parameter λ , but still the different loop functions for t and t' -quarks prevent us from deriving simple analytical expressions as compared to the decoupling scenarios considered in [31].

3 Numerical Procedure and Results

In this section we are going to present the results of our numerical analysis. First we will discuss the general procedure we have used. Subsequently we will discuss our results for various observables recalling, where necessary, some definitions from our previous papers on $D^0 - \bar{D}^0$ mixing. For all plots we will use, if not indicated otherwise, the same colour-coding as in our previous analysis of the SM4 [20] where we dealt with mixing and rare decays in the K and B meson systems. This colour coding is given in Table 1 and allows to distinguish between sets of SM4 parameters for which the CP asymmetry $S_{\psi\phi}(B_s)$ in the B_s system and the branching ratio $\text{Br}(B_s \rightarrow \mu^+ \mu^-)$ assume certain values. Furthermore, the *light blue* points correspond to $\text{Br}(K_L \rightarrow \pi^0 \nu \bar{\nu}) > 2 \cdot 10^{-10}$, while *dark blue* points indicate $\text{Br}(K_L \rightarrow \pi^0 \nu \bar{\nu}) \leq 2 \cdot 10^{-10}$. This colour coding will give us some insight into the correlations between the different mesonic systems.

3.1 Strategy for Our Phenomenological Analysis

For our analysis we use the points generated for our previous analysis [20]. For the LD contributions Γ_{12}^{LD} and M_{12}^{LD} we adopt the same procedure as in [33]: We scan flatly over

	BS1 (yellow)	BS2 (green)	BS3 (red)
$S_{\psi\phi}(B_s)$	0.04 ± 0.01	0.04 ± 0.01	≥ 0.4
$\text{Br}(B_s \rightarrow \mu^+ \mu^-)$	$(2 \pm 0.2) \cdot 10^{-9}$	$(3.2 \pm 0.2) \cdot 10^{-9}$	$\geq 6 \cdot 10^{-9}$

Table 1: Three scenarios for $S_{\psi\phi}(B_s)$ and $\text{Br}(B_s \rightarrow \mu^+ \mu^-)$.

parameter	value	parameter	value
η_{cc}	1.51 ± 0.24 [46]	m_D	$(1.86484 \pm 0.00017)\text{GeV}$
η_{tt}	0.5765 ± 0.0065 [40]	$\bar{\tau}_D$	$(0.4101 \pm 0.0015)\text{ps}$
η_{ct}	0.47 ± 0.04 [47]		
F_D	$(0.212 \pm 0.014) \text{ GeV}$ [48]	$m_c(m_c)$	$(1.268 \pm 0.009) \text{ GeV}$ [49, 50]
$\hat{B}_D$	$1.18^{+0.07}_{-0.05}$	$m_b(m_b)$	$(4.20^{+0.17}_{-0.07}) \text{ GeV}$ [43]

Table 2: Values of the input parameters used in our analysis.

the intervals

$$-0.02 \text{ ps}^{-1} \leq M_{12}^{\text{LD}} \leq 0.02 \text{ ps}^{-1}, \quad (3.15)$$

$$-0.04 \text{ ps}^{-1} \leq \Gamma_{12}^{\text{LD}} \leq 0.04 \text{ ps}^{-1}, \quad (3.16)$$

while requiring x_D , y_D and $|q/p|$ to be within 2σ of their experimental values. The latter with their 1σ ranges are given by [43–45]

$$\begin{aligned}
x_D &= (0.98^{+0.24}_{-0.26}) \%, & y_D &= (0.83 \pm 0.16) \%, \\
|q/p| &= (0.87^{+0.17}_{-0.15}), & \varphi &= (-8.5^{+7.4}_{-7.0})^\circ, \\
\eta_f S_f(D) &= (-0.248 \pm 0.496) \%,
\end{aligned} \quad (3.17)$$

with φ being the phase of q/p and the asymmetry $\eta_f S_f(D)$ defined in (3.20). BaBar recently presented preliminary results for x_D compatible with $x_D = 0$. Because of the large relative error and the presence of potentially large LD contributions, we will however not elaborate on these new results and use the HFAG averages.

3.2 The SD Contribution to $D^0 - \bar{D}^0$ Mixing

We now turn our attention to the SD contributions which are summarised by the amplitude M_{12}^D in (2.10). In Fig. 1 we show this amplitude in the complex plane. The striking features of this plot are the following

- i) We observe a clustering of the *red* points (indicating large $S_{\psi\phi}(B_s)$ and $\text{Br}(B_s \rightarrow \mu^+\mu^-)$) along the $\text{Im}(M_{12}^D) = 0$ axis. Furthermore, for this class of points, significant positive values of $\text{Im}(M_{12}^D)$ are possible, while a rather stringent lower bound, $\text{Im}(M_{12}^D) \gtrsim -2 \cdot 10^{-3}$ is found.
- ii) For $\text{Re}(M_{12}^D) > 0$, values ranging from very small up to a factor of four larger than the maximal values of $\text{Im}(M_{12}^D)$ are possible. For $\text{Re}(M_{12}^D) < 0$, only very small values are allowed due to the stringent ε_K constraint.
- iii) The largest values for $\text{Im}(M_{12}^D)$ are found in the *light blue* scenario corresponding to large values of $\text{Br}(K_L \rightarrow \pi^0\nu\bar{\nu})$, which is governed by CP violation in the K system. We conclude that large CP violating effects in the D system entail large CP violating effects also in the $K \rightarrow \pi\nu\bar{\nu}$ system, while the converse is not necessarily true.




Figure 1: The imaginary part as a function of the real part of M_{12}^D .

3.3 CP Violation in the $D^0 - \bar{D}^0$ System

In Fig. 2 we show the semileptonic CP asymmetry $a_{\text{SL}}(D)$ as a function of $|q/p|$. This asymmetry represents CP violation in $\mathcal{L}(\Delta C = 2)$ and is related to $|q/p|$ simply by

$$a_{\text{SL}}(D) \equiv \frac{\Gamma(D^0(t) \rightarrow \ell^- \bar{\nu} K^{+(*)}) - \Gamma(\bar{D}^0 \rightarrow \ell^+ \nu K^{-(*)})}{\Gamma(D^0(t) \rightarrow \ell^- \bar{\nu} K^{+(*)}) + \Gamma(\bar{D}^0 \rightarrow \ell^+ \nu K^{-(*)})} = \frac{|q|^4 - |p|^4}{|q|^4 + |p|^4} \approx 2 \left(\left| \frac{q}{p} \right| - 1 \right). \quad (3.18)$$




Figure 2: The semileptonic CP asymmetry $a_{\text{SL}}(D)$ as a function of $|q/p|$.

In writing the last expression, we have assumed that $|q/p| - 1$ is much smaller than unity, which is supported by the data listed in (3.17).

We observe—in accordance with Fig. 1—that in the *red* scenario, in which CP violation in the B_s system is large, the asymmetry $a_{\text{SL}}(D)$ in the D system in the allowed range of $|q/p|$ is small but still much larger than in the SM3:

$$[a_{\text{SL}}(D)]_{\text{SM3}} \approx 1 \cdot 10^{-4}. \quad (3.19)$$

Significantly larger values are found in the other scenarios in which $S_{\psi\phi}(B_s)$ is SM-like but $\text{Br}(K_L \rightarrow \pi^0 \nu \bar{\nu})$ might be strongly enhanced.

Of particular interest is the time-dependent CP asymmetry S_f defined by [32]

$$\frac{\Gamma(D^0(t) \rightarrow f) - \Gamma(\bar{D}^0(t) \rightarrow f)}{\Gamma(D^0(t) \rightarrow f) + \Gamma(\bar{D}^0(t) \rightarrow f)} \equiv S_f(D) \frac{t}{2\tau_D}, \quad (3.20)$$

which is given by

$$\eta_f S_f(D) \simeq - \left[y_D \left(\left| \frac{q}{p} \right| - \left| \frac{p}{q} \right| \right) \cos \varphi - x_D \left(\left| \frac{q}{p} \right| + \left| \frac{p}{q} \right| \right) \sin \varphi \right], \quad (3.21)$$

where $\eta_f = \pm 1$ is the CP parity of the final state f . The SM3 prediction for $\eta_f S_f(D)$ is

$$[\eta_f S_f(D)]_{\text{SM3}} \approx -2 \cdot 10^{-6}. \quad (3.22)$$

Other useful related formulae can be found in [31–33, 36].

In the absence of significant CP phases in the decay amplitudes for $D^0 \rightarrow f$, the asymmetry S_f displays a strong correlation with the semileptonic asymmetry $a_{\text{SL}}(D)$ [32],

$$\eta_f S_f(D) \simeq - \frac{x_D^2 + y_D^2}{y_D} a_{\text{SL}}(D). \quad (3.23)$$

A similar correlation is familiar from the B_s system [51–53]. In the presence of significant phases in decay amplitudes $D^0 \rightarrow f$, formula (4.23) in [32] instead of (3.21, 3.23) applies.

In Fig. 3 we show $\eta_f S_f(D)$ as a function of the phase φ (left panel) and as a function of $a_{\text{SL}}(D)$ (right panel). These plots are familiar from the model-independent analysis in [33] and are self-explanatory. In particular in the *red* scenario the points cluster around $\eta_f S_f(D) \approx 0$, but still $\eta_f S_f(D)$ can be much larger than found in the SM3. Again extreme values are only found for the *light blue* points corresponding to a large enhancement of $\text{Br}(K_L \rightarrow \pi^0 \nu \bar{\nu})$.




Figure 3: Model independent correlations between $\eta_f S_f(D)$ and φ (left panel) and a_{SL} (right panel).

3.4 Correlation of $D^0 - \bar{D}^0$ Mixing with the K and B Systems

In order to get some insight into the correlations of the D system with other meson systems, we show in Figs. 4–7 a symphony of plots which depict the dependence of $\eta_f S_f(D)$ on a number of prominent observables in the K and $B_{s,d}$ systems:

- In the left and right panels of Fig. 4 we plot $\eta_f S_f(D)$ as a function of $\text{Br}(K_L \rightarrow \pi^0 \nu \bar{\nu})$ and $\text{Br}(K^+ \rightarrow \pi^+ \nu \bar{\nu})$. We observe that values of $|\eta_f S_f(D)|$ up to $\sim 1\%$ are found in the region where $\text{Br}(K_L \rightarrow \pi^0 \nu \bar{\nu})$ and $\text{Br}(K^+ \rightarrow \pi^+ \nu \bar{\nu})$ are strongly enhanced, but also $\eta_f S_f(D) = 0$ is possible in this region.
- In the left and right panels of Fig. 5 we plot $\eta_f S_f(D)$ as a function of $\text{Br}(B_d \rightarrow \mu^+ \mu^-)$ and $\text{Br}(B_s \rightarrow \mu^+ \mu^-)$, respectively. As expected, for very large values of $\text{Br}(B_s \rightarrow \mu^+ \mu^-)$, $\eta_f S_f(D)$ tends to be within the experimental 1σ range and thus small. On the other hand, we find that even for SM-like values of $B_s \rightarrow \mu\mu$ and $S_{\psi\phi}(B_s)$, maximal values in $\eta_f S_f(D)$ are attainable. Both effects are even more pronounced in the case of $\text{Br}(B_d \rightarrow \mu^+ \mu^-)$. Thus for some values of $\text{Br}(B_d \rightarrow$

$\mu^+\mu^-$) in conjunction with $\eta_f S_f(D)$ we are able to predict ranges for $\text{Br}(K_L \rightarrow \pi^0 \nu \bar{\nu})$. The D system in this sense acts as a link between the K and B systems.

- In Fig. 6 we plot $\eta_f S_f(D)$ as a function of the mixing-induced CP asymmetry $S_{\psi\phi}(B_s)$ in the B_s system. While many of the features in this plot are self-explanatory on the basis of previous plots, the striking result is that for $S_{\psi\phi}(B_s) > 0$ as signalled by CDF and D0 data, $\eta_f S_f(D)$ is predicted to be within its experimental 1σ range and thus small.
- As was pointed out in [20], the measured value of ε'/ε practically eliminates large CP violating effects in the B_s system. In order to study this constraint in the D system, in Fig. 7 we show ε'/ε as a function of $\eta_f S_f(D)$ for different values of the non-perturbative parameters (R_6, R_8). We observe that ε'/ε additionally implies *negative* or *small positive* values for $\eta_f S_f(D)$. Large negative values for $\eta_f S_f(D)$ however are only possible if also significant CP violation from 4G in the $K \rightarrow \pi \nu \bar{\nu}$ system is present.




Figure 4: $\eta_f S_f(D)$ as a function of $\text{Br}(K^+ \rightarrow \pi^+ \nu \bar{\nu})$ (left panel) and $\text{Br}(K_L \rightarrow \pi^0 \nu \bar{\nu})$ (right panel).




Figure 5: $\eta_f S_f(D)$ as a function of $\text{Br}(B_d \rightarrow \mu^+ \mu^-)$ (left panel) and $\text{Br}(B_s \rightarrow \mu^+ \mu^-)$ (right panel).




Figure 6: $\eta_f S_f(D)$ as a function of $S_{\psi\phi}(B_s)$.




Figure 7: ε'/ε as a function of the CP asymmetry $\eta_f S_f(D)$ for four different scenarios of the non-perturbative parameters: $(R_6, R_8) = (1.0, 1.0)$ (upper left panel), $(1.5, 0.8)$ (upper right panel), $(2.0, 1.0)$ (lower left panel) and $(1.5, 0.5)$ (lower right panel).

3.5 Interplay between CP Violation in the K , D and B_s Systems

Let us, for illustration, consider an extreme situation where we take the constraints on the 4G parameters from the present experimental and theoretical situation regarding ε'/ε and $S_{\psi\phi}(B_s)$ at face value. To this end, we restrict our set of allowed 4G parameter values to fulfil the 2σ range for ε'/ε and either $S_{\psi\phi}(B_s) > 0$ or $S_{\psi\phi}(B_s) > 0.2$. For the remaining points, we plot again $\eta_f S_f(D)$ as a function of $a_{\text{SL}}(D)$ and thus obtain a rather constrained prediction for the impact of the 4G in the charm system, as shown in Fig. 8. In particular, we find that the requirement $S_{\psi\phi}(B_s) > 0$ in conjunction with the imposition of the constraint from ε'/ε drastically reduces the possible NP effects with respect to those shown in Fig. 3, but still allows for moderate effects in $a_{\text{SL}}(D)$ and $\eta_f S_f(D)$, where $\eta_f S_f(D) < [\eta_f S_f(D)]_{\text{SM3}}$ is slightly favoured. Demanding $S_{\psi\phi}(B_s) > 0.2$ on the other hand precludes almost any deviation from the SM3 prediction, regardless of the values of the hadronic parameters R_6 , R_8 entering ε'/ε .




Figure 8: Prediction for the CP violating parameters $\eta_f S_f(D)$ vs. $a_{\text{SL}}(D)$ in the charm system for 4G parameters restricted to fulfil the 2σ range for ε'/ε and $S_{\psi\phi}(B_s) > 0$ (left panel) or $S_{\psi\phi}(B_s) > 0.2$ (right panel). The colour coding is according to Table 5 in [20] and indicates the particular values of the hadronic parameters R_6 and R_8 entering ε'/ε .

3.6 Scaling Scenarios

In [20], we have shown that it is useful to characterise the hierarchical structures in the 4G mixing matrix by dividing the parameter space into subclasses, which are defined by the scaling of the 4G mixing angles with the Wolfenstein parameter λ and the correlation between the 4G CP phases δ_{14} and δ_{24} which is induced by the constraints from precision flavour observables. Let us, as an example, investigate the correlation between the time-

dependent CP asymmetry $\eta_f S_f(D)$ and the semileptonic asymmetry $a_{\text{SL}}(D)$ in the D^0 decays for the individual subclasses defined in [20], see Fig. 9. (We have not shown results for subclasses 4 and 5, as they only lead to tiny deviations from the SM, since condition (2.14) is not fulfilled.) As in the case of B -meson and K -meson observables, the correlations between $a_{\text{SL}}(D)$ and $\eta_f S_f(D)$ are quite different for different sub-classes, allowing again for a clear separation with respect to the possible scaling behaviour of the 4G mixing angles. Similar plots can be drawn for the other correlations studies above.

As a further example we study the 4G parameter points identified in Section 3.5 that are consistent with both the recent experimental data from CDF and D0 implying sizable $S_{\psi\phi}(B_s)$, and the measurement of ε'/ε . When studying the scaling of the 4G mixing angles and the correlation between the 4G phases for these points, we find—for example requiring that $S_{\psi\phi}(B_s) > 0.2$ —that the bulk of parameter values has $(\theta_{14}, \theta_{24}, \theta_{34}) \sim (\lambda^3, \lambda^2, \lambda)$ and $-\pi < \delta_{14} \simeq \delta_{24} < 0$. In terms of the subclasses which we have discussed above, this corresponds to class 1a (and to a somewhat lesser extent class 3b). A few points with $(\theta_{14}, \theta_{24}, \theta_{34}) \sim (\lambda^4, \lambda^3, \lambda)$ survive, too, but as we already explained above, they correspond to the uninteresting case where the 4G effects in the charm system are very small. Thus, as was the case in our previous analysis [20], we find that there is a strong connection between the phenomenological predictions of the SM4 on the one hand and the scaling of the mixing parameters θ_{i4} in powers of the Wolfenstein parameter λ on the other hand.




Figure 9: Correlation between $\eta_f S_f(D)$ and $a_{SL}(D)$ for the individual subclasses defined in [20]. The colour coding of the points is as explained in Section 3.

4 Conclusions

In the present paper we have analysed the impact of a 4G of quarks on the $D^0 - \bar{D}^0$ system taking into account all existing constraints. In addition we have investigated the correlations of the D system with the prominent observables in the K and $B_{d,s}$ systems. The main messages from our analysis are as follows:

- Large 4G effects in the CP violating observables $a_{\text{SL}}(D)$ and $\eta_f S_f(D)$ are possible and consistent with the available data. The effects are found to be as large as a few times $\mathcal{O}(10^{-1})$ for $a_{\text{SL}}(D)$ and up to $\sim 1.5\%$ for $\eta_f S_f(D)$, although this seems to be disfavoured by a small $\varphi_{\text{exp}} \sim -8^\circ$. Compared to the SM3 predictions for these asymmetries, $a_{\text{SL}}(D) \sim 10^{-4}$ and $\eta_f S_f(D) \sim 10^{-6}$, even the more moderate enhancements allowed by the current data are still spectacular.
- The large 4G effects in the K , B_d and B_s systems found by us in [20] are still compatible with the current constraints from the charm data.
- While simultaneous large 4G effects in the K and D systems are possible, large effects in B_d generally disfavour large NP effects in the D system. In the B_s system large CP *conserving* effects are found to be possible regardless of the predictions for the D system, while large CP *violating* effects can only occur if $\eta_f S_f(D)$ does not deviate from the experimental measurement by much more than 1σ . Conversely, significant enhancement of $S_{\psi\phi}(B_s)$ above the SM3 value will not allow large CP violating effects in the D system with the 4G scenario.
- Additional imposition of the ε'/ε constraint significantly diminishes 4G effects in CP violating observables in the D system. This observation is to a large extent independent of the actual values of the hadronic parameters R_6 and R_8 entering ε'/ε that still suffer from significant theoretical uncertainties.

In the light of these findings we are looking forward to improved data on charm and its correlations with B and K physics measurements to be discovered in the upcoming decade.

Acknowledgements

We thank A. Kronfeld for the illuminating discussion related to the hadronic parameter $\hat{B}_D$. AJB, BD, TF and TH want to thank the Galileo Galilei Institute for Theoretical Physics for their hospitality. AJB and TF thank the INFN for partial support during the initial steps of this work. This research was partially supported by the Cluster of

Excellence ‘Origin and Structure of the Universe’, the Graduiertenkolleg GRK 1054 of DFG and by the German ‘Bundesministerium für Bildung und Forschung’ under contract 05H09WOE.

References

- [1] P. H. Frampton, P. Q. Hung, and M. Sher, *Quarks and leptons beyond the third generation*, *Phys. Rept.* **330** (2000) 263, hep-ph/9903387 .
- [2] B. Holdom *et al.*, *Four Statements about the Fourth Generation*, 0904.4698.
- [3] G. D. Kribs, T. Plehn, M. Spannowsky, and T. M. P. Tait, *Four generations and Higgs physics*, *Phys. Rev.* **D76** (2007) 075016, 0706.3718 .
- [4] M. Hashimoto, *Constraints on Mass Spectrum of Fourth Generation Fermions and Higgs Bosons*, 1001.4335.
- [5] J. Alwall *et al.*, *Is $V_{tb} \simeq 1$?*, *Eur. Phys. J.* **C49** (2007) 791–801, hep-ph/0607115 .
- [6] M. S. Chanowitz, *Bounding CKM Mixing with a Fourth Family*, *Phys. Rev.* **D79** (2009) 113008, 0904.3570 .
- [7] V. A. Novikov, A. N. Rozanov, and M. I. Vysotsky, *Once more on extra quark-lepton generations and precision measurements*, 0904.4570.
- [8] J. Erler and P. Langacker, *Precision Constraints on Extra Fermion Generations*, 1003.3211.
- [9] P. Q. Hung and C. Xiong, *Renormalization Group Fixed Point with a Fourth Generation: Higgs-induced Bound States and Condensates*, 0911.3890.
- [10] P. Q. Hung and C. Xiong, *Renormalization Group Fixed Point with a Fourth Generation: Solution to the hierarchy problem*, 0911.3892.
- [11] K. S. Babu, X. G. He, X. Li, and S. Pakvasa, *FOURTH GENERATION SIGNATURES IN D^0 - $\bar{D}^0$ MIXING AND RARE D DECAYS*, *Phys. Lett.* **B205** (1988) 540.
- [12] A. Arhrib and W.-S. Hou, *Effect of fourth generation CP phase on $b \rightarrow s$ transitions*, *Eur. Phys. J.* **C27** (2003) 555–561, hep-ph/0211267 .

- [13] W.-S. Hou, M. Nagashima, and A. Soddu, *Enhanced $K_L \rightarrow \pi^0 \nu \bar{\nu}$ from direct CP violation in $B \rightarrow K\pi$ with four generations*, *Phys. Rev.* **D72** (2005) 115007, hep-ph/0508237 .
- [14] W.-S. Hou, M. Nagashima, and A. Soddu, *Large time-dependent CP violation in B_s^0 system and finite $D^0 - \bar{D}^0$ mass difference in four generation standard model*, *Phys. Rev.* **D76** (2007) 016004, hep-ph/0610385 .
- [15] A. Soni, A. K. Alok, A. Giri, R. Mohanta, and S. Nandi, *The Fourth family: A Natural explanation for the observed pattern of anomalies in B -CP asymmetries*, 0807.1971.
- [16] J. A. Herrera, R. H. Benavides, and W. A. Ponce, *Flavor changing neutral currents with a fourth family of quarks*, *Phys. Rev.* **D78** (2008) 073008, 0810.3871 .
- [17] M. Bobrowski, A. Lenz, J. Riedl, and J. Rohrwild, *How much space is left for a new family of fermions?*, *Phys. Rev.* **D79** (2009) 113006, 0902.4883 .
- [18] G. Eilam, B. Melic, and J. Trampetic, *CP violation and the 4th generation*, *Phys. Rev.* **D80** (2009) 116003, 0909.3227 .
- [19] A. Soni, A. K. Alok, A. Giri, R. Mohanta, and S. Nandi, *SM with four generations: Selected implications for rare B and K decays*, 1002.0595.
- [20] A. J. Buras, B. Duling, T. Feldmann, T. Heidsieck, C. Promberger, and S. Recksiegel, *Patterns of Flavour Violation in the Presence of a Fourth Generation of Quarks and Leptons*, 1002.2126.
- [21] W. S. Hou and C. Y. Ma, *Flavor and CP Violation with Fourth Generations Revisited*, 1004.2186.
- [22] Z. Murdock, S. Nandi, and Z. Tavartkiladze, *Perturbativity and a Fourth Generation in the MSSM*, *Phys. Lett.* **B668** (2008) 303–307, 0806.2064 .
- [23] R. M. Godbole, S. K. Vempati, and A. Wingerter, *Four Generations: SUSY and SUSY Breaking*, 0911.1882.
- [24] M. Blanke *et al.*, *Particle antiparticle mixing, ϵ_K , $\Delta(\Gamma(q))$, $A_{SL}(q)$, $A_{CP}(B_d \rightarrow \psi K_S)$, $A_{CP}(B_s \rightarrow \psi \phi)$ and $B \rightarrow X_S, d\gamma$ in the littlest Higgs model with T -parity*, *JHEP* **12** (2006) 003, hep-ph/0605214 .
- [25] M. Blanke *et al.*, *Rare and CP-Violating K and B Decays in the Littlest Higgs Model with T -Parity*, *JHEP* **01** (2007) 066, hep-ph/0610298 .

- [26] M. Blanke, A. J. Buras, B. Duling, S. Recksiegel, and C. Tarantino, *FCNC Processes in the Littlest Higgs Model with T-Parity: a 2009 Look*, *Acta Phys. Polon. B* **41** (2010) 657, 0906.5454 .
- [27] M. Blanke, A. J. Buras, B. Duling, S. Gori, and A. Weiler, *$\Delta F = 2$ Observables and Fine-Tuning in a Warped Extra Dimension with Custodial Protection*, *JHEP* **03** (2009) 001, 0809.1073 .
- [28] M. Blanke, A. J. Buras, B. Duling, K. Gemmler, and S. Gori, *Rare K and B Decays in a Warped Extra Dimension with Custodial Protection*, *JHEP* **03** (2009) 108, 0812.3803 .
- [29] W. Altmannshofer, A. J. Buras, S. Gori, P. Paradisi, and D. M. Straub, *Anatomy and Phenomenology of FCNC and CPV Effects in SUSY Theories*, *Nucl. Phys.* **B830** (2010) 17–94, 0909.1333 .
- [30] M. Bobrowski, A. Lenz, J. Riedl, and J. Rohrwild, *How large can the SM contribution to CP violation in $D^0 - \bar{D}^0$ mixing be?*, 1002.4794.
- [31] K. Blum, Y. Grossman, Y. Nir, and G. Perez, *Combining $K - \bar{K}$ mixing and $D - \bar{D}$ mixing to constrain the flavor structure of new physics*, *Phys. Rev. Lett.* **102** (2009) 211802, 0903.2118 .
- [32] I. I. Bigi, M. Blanke, A. J. Buras, and S. Recksiegel, *CP Violation in $D^0 - \bar{D}^0$ Oscillations: General Considerations and Applications to the Littlest Higgs Model with T-Parity*, *JHEP* **07** (2009) 097, 0904.1545 .
- [33] W. Altmannshofer, A. J. Buras, and P. Paradisi, *A Lower Bound on hadronic EDMs from CP Violation in $D^0 - \bar{D}^0$ mixing in SUSY Alignment Models*, *Phys. Lett.* **B688** (2010), no. 2-3 202, 1001.3835 .
- [34] E. Golowich, J. Hewett, S. Pakvasa, and A. A. Petrov, *Implications of $D^0 - \bar{D}^0$ Mixing for New Physics*, *Phys. Rev.* **D76** (2007) 095009, 0705.3650 .
- [35] **Particle Data Group** Collaboration, C. Amsler *et al.*, *Review of particle physics*, *Phys. Lett.* **B667** (2008) 1.
- [36] A. L. Kagan and M. D. Sokoloff, *On Indirect CP Violation and Implications for $D^0 - \bar{D}^0$ and $B_s - \bar{B}_s$ mixing*, *Phys. Rev.* **D80** (2009) 076008, 0907.3917 .
- [37] R. Gupta, T. Bhattacharya, and S. R. Sharpe, *Matrix elements of 4-fermion operators with quenched Wilson fermions*, *Phys. Rev.* **D55** (1997) 4036–4054, hep-lat/9611023 .

- [38] **UKQCD** Collaboration, L. Lellouch and C. J. D. Lin, *Standard model matrix elements for neutral B meson mixing and associated decay constants*, *Phys. Rev. D* **D64** (2001) 094501, [hep-ph/0011086](#) .
- [39] H.-W. Lin, S. Ohta, A. Soni, and N. Yamada, *Charm as a domain wall fermion in quenched lattice QCD*, *Phys. Rev. D* **D74** (2006) 114506, [hep-lat/0607035](#) .
- [40] A. J. Buras, M. Jamin, and P. H. Weisz, *Leading and next-to-leading QCD corrections to ϵ parameter and $B^0 - \bar{B}^0$ mixing in the presence of a heavy quark*, *Nucl. Phys. B* **B347** (1990) 491–536.
- [41] S. Bethke, *The 2009 World Average of $\alpha_s(M_Z)$* , *Eur. Phys. J. C* **C64** (2009) 689–703, [0908.1135](#) .
- [42] M. Blanke, A. J. Buras, S. Recksiegel, and C. Tarantino, *The Littlest Higgs Model with T-Parity Facing CP-Violation in $B_s - \bar{B}_s$ Mixing*, [0805.4393](#).
- [43] **Heavy Flavor Averaging Group (HFAG)** Collaboration, E. Barberio *et al.*, *Averages of b -hadron properties at the end of 2006*, [0704.3575](#). Updates available on <http://www.slac.stanford.edu/xorg/hfag>.
- [44] **Heavy Flavor Averaging Group** Collaboration, E. Barberio *et al.*, *Averages of b -hadron and c -hadron Properties at the End of 2007*, [0808.1297](#).
- [45] A. J. Schwartz, *$D^0 - \bar{D}^0$ Mixing and CP Violation: HFAG Combination of Parameters*, [0911.1464](#).
- [46] S. Herrlich and U. Nierste, *Enhancement of the $K_L - K_S$ mass difference by short distance QCD corrections beyond leading logarithms*, *Nucl. Phys. B* **B419** (1994) 292–322, [hep-ph/9310311](#) .
- [47] S. Herrlich and U. Nierste, *Indirect CP violation in the neutral kaon system beyond leading logarithms*, *Phys. Rev. D* **D52** (1995) 6505–6518, [hep-ph/9507262](#) .
- [48] V. Lubicz and C. Tarantino, *Flavour physics and Lattice QCD: averages of lattice inputs for the Unitarity Triangle Analysis*, *Nuovo Cim.* **123B** (2008) 674–688, [0807.4605](#) .
- [49] J. Laiho, R. S. Van de Water, and E. Lunghi, *Lattice QCD inputs to the CKM unitarity triangle analysis*, [0910.2928](#).
- [50] **HPQCD** Collaboration, I. Allison *et al.*, *High-Precision Charm-Quark Mass from Current-Current Correlators in Lattice and Continuum QCD*, *Phys. Rev. D* **D78** (2008) 054513, [0805.2999](#) .

- [51] Z. Ligeti, M. Papucci, and G. Perez, *Implications of the measurement of the $B^0_s - \bar{B}^0_s$ mass difference*, *Phys. Rev. Lett* **97** (2006) 101801, [hep-ph/0604112](#) .
- [52] M. Blanke, A. J. Buras, D. Guadagnoli, and C. Tarantino, *Minimal Flavour Violation Waiting for Precise Measurements of ΔM_s , $S_{\psi\phi}$, A^s_{SL} , $|V_{ub}|$, γ and $B^0_{s,d} \rightarrow \mu^+\mu^-$* , *JHEP* **10** (2006) 003, [hep-ph/0604057](#) .
- [53] Y. Grossman, Y. Nir, and G. Perez, *Testing New Indirect CP Violation*, *Phys. Rev. Lett.* **103** (2009) 071602, [0904.0305](#) .